Your heart failure medicines tracker

According to the American College of Cardiology (ACC) and the American Heart Association (AHA), these are the different types of medications used to treat heart failure. Your doctor may have you on one or more of these medicines. To help you stay on track:



- 1. Check off which medications you're on
- 2. Write down the current dose

Bring this tracker with you to your doctor's appointments. Discuss any side effects or symptoms you may be having. This will help you and your doctor figure out which meds are right for you and what changes (if any) need to be made. Try for ONE positive change at each doctor's visit.

		Generic Medicine Name (Brand Name)	Current Dose	Target Dose
Potr	ta-blockers	bisoprolol (Zebeta)	mg/day	10 mg/day
		metoprolol succinate (Toprol)	mg/day	200 mg/day
Dec		carvedilol (Coreg)	mg (2 times/day)	25-50 mg (2 times/day)
Cor Enz	giotensin- inverting zyme (ACE) nibitors	captopril (Capoten)	mg (3 times/day)	50 mg (3 times/day)
		ramipril (Altace)	mg/day	10 mg/day
		enalapril (Vasotec)	mg/day	10-20 mg/day
		lisinopril (Prinivil, Zestril)	mg/day	20-40 mg/day
Ang	giotensin II ceptor ockers (ARBs)	candesartan (Atacand)	mg/day	32 mg/day
Rec		Iosartan (Cozaar)	mg/day	150 mg/day
		valsartan (Diovan)	mg (2 times/day)	160 mg (2 times/day)
Rec Nep	giotensin eptor- orilysin bitors (ARNIs)	sacubitril/valsartan (Entresto)	mg (2 times/day)	97/103 mg (2 times/day)
Rec Ant	losterone ceptor tagonist RAs)	eplerenone (Inspra)	mg/day	50 mg/day
		spironolactone (Aldactone)	mg/day	25-50 mg/day



Generic Medicine Name (Brand Name)	Current Dose	Target Dose
hydralazine (Apresoline)	mg (3 times/day)	75 mg (3 times/day)
isosorbide dinitrate (Angitak)	mg (3 times/day)	40 mg (3 times/day)
Fixed-dose hydralazine/isosorbide dinitrate	tablets (3 times/day)	2 tablets (3 times/day)
empagliflozin (Jardiance)	mg/day	10 mg/day
dapagliflozin (Farxiga)	mg/day	10 mg/day
canagliflozin (Invokana)	mg/day	100 mg/day
ivabradine (Corlanor)	mg (2 times/day)	Titrate to heart rate 50-60/min
	mig (2 times/day)	Max Dose 7.5 mg (2 times/day)

 $Yancy, et al.\ 2017\ ACC\ Expert\ Consensus\ Decision\ Pathway\ for\ Optimization\ of\ Heart\ Failure\ Treatment:\ Answers\ to\ 10\ Pivotal\ Issues\ About\ Heart\ Failure\ With\ Reduced\ Ejection\ Fraction:\ A\ Report\ of\ the\ American\ College\ of\ Cardiology\ Task\ Force\ on\ Expert\ Consensus\ Decision\ Pathways.$

Yancy, et al. 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines.

Murphy et al. Heart Failure With Reduced Ejection Fraction: A Review. JAMA 2020.

Note: Target dose is the optimal dose or maximum tolerated dose.

Vasodilators

Sodium-Glucose Co-transporter 2 Inhibitors (SGLT2)

Sinoatrial Node Inhibitor

Check off any symptoms or issues that you're currently having:

Dizzy Short of breath when Change in taste leaning over (bendopnea) Tired Dry mouth A rapid or irregular Dry cough Vision issues (bright lights) heartbeat Short of breath Cold hands or feet Need more than two pillows to sleep Chest pain Depressed (need to be propped up) Swelling Other: Wake up suddenly at night with Headaches trouble breathing

Cough produces

mucus or pink sputum



Lack of appetite

Early satiety or

feeling full early